

Fuzzy Anti-norm and Fuzzy α -anti-convergence

Bivas Dinda, T.K. Samanta and Iqbal H. Jebril

Department of Mathematics,
Mahishamuri Ramkrishna Vidyapith, West Bengal, India.

e-mail: bvsdinda@gmail.com

Department of Mathematics, Uluberia College, West Bengal, India.

e-mail: mumpu_tapas5@yahoo.co.in

Department of Mathematics, King Faisal University, Saudi Arabia.

e-mail: iqbal501@hotmail.com

Abstract

Fuzzy anti-norm and corresponding α -norms are defined. A few properties of finite dimensional fuzzy anti-normed linear space are studied. Fuzzy α -anti-convergence and fuzzy α -anti-complete linear space are defined and a few of their properties are studied.

Keywords: *Fuzzy anti-norm, α -norm, Fuzzy α -anti-convergence, fuzzy α -anti-complete.*

2010 Mathematics Subject Classification: 03E72, 46S40.

1 Introduction

The theory of fuzzy sets was first introduced by Zadeh[12] in 1965. Thereafter the concept of fuzzy norm was introduced by Katsaras [8] in 1984. In 1992, Felbin[6] introduced the idea of fuzzy norm on a linear space. Cheng-Moderson [4] introduced another idea of fuzzy norm on a linear space whose associated metric is same as the associated metric of Kramosil-Michalek [10]. Bag and Samanta[2] modified the definition of fuzzy norm of Cheng-Moderson[4] and then studied different properties of fuzzy normed linear space..

In this paper, the definition of fuzzy anti-norm on a linear space introduced by Jebril and Samanta[1], has been modified and then the Riesz lemma and a few important properties of finite dimensional fuzzy anti-normed linear space has been established. We define fuzzy α -anti-convergence, fuzzy α -anti-cauchy sequence, fuzzy α -anti-completeness and relations among them are studied.

2 Preliminaries

This section contains a few basic definitions and preliminary results which will be needed in the sequel.

Definition 2.1 [11]. A binary operation $\diamond : [0, 1] \times [0, 1] \longrightarrow [0, 1]$ is a t -conorm if $\diamond$ satisfies the following conditions :

- (i) $\diamond$ is commutative and associative ,
- (ii) $a \diamond 0 = a \quad \forall a \in [0, 1]$,
- (iii) $a \diamond b \leq c \diamond d$ whenever $a \leq c, b \leq d$ and $a, b, c, d \in [0, 1]$.

Remark 2.2 [9]. (a) For any $r_1, r_2 \in (0, 1)$ with $r_1 > r_2$, there exist $r_3 \in (0, 1)$ such that $r_1 > r_3 \diamond r_2$.

(b) For any $r_4 \in (0, 1)$, there exist $r_5 \in (0, 1)$ such that and $r_5 \diamond r_5 \leq r_4$.

Definition 2.3 [1]. Let U be a linear space over the real field F . A fuzzy subset N^* of $U \times R$ such that for all $x, u \in U$ and $c \in F$

- (N*1) For all $t \in R$ with $t \leq 0$, $N^*(x, t) = 1$;
- (N*2) For all $t \in R$ with $t > 0$, $N^*(x, t) = 0$ if and only if $x = \theta$;
- (N*3) For all $t \in R$ with $t > 0$, $N^*(cx, t) = N^*(x, \frac{t}{|c|})$ if $c \neq 0, c \in F$;
- (N*4) For all $s, t \in R$ with $N^*(x+u, s+t) \leq \max \{ N^*(x, s), N^*(y, t) \}$;
- (N*5) $N^*(x, t)$ is a non-increasing function of $t \in R$ and $\lim_{t \rightarrow \infty} N^*(x, t) = 0$.

Then (U, N^*) is called fuzzy antinormed linear space. Further it is assumed

- (N*6) For all $t \in R$ with $t > 0$, $N^*(x, t) < 1$ implies $x = \theta$.

Definition 2.4 [1]. Let N^* be a fuzzy antinorm on U satisfying (N*6). Define

$$\|x\|_{\alpha}^* = \inf \{ t > 0 : N^*(x, t) < \alpha \}$$

Theorem 2.5 [1]. Let (U, N^*) be a fuzzy antinormed linear space. Then $\{\|\cdot\|_{\alpha}^* : \alpha \in (0, 1)\}$ is a decreasing family of norms on U .

Definition 2.6 [1]. Let (U, N^*) be a fuzzy antinormed linear space. A sequence $\{x_n\}_n$ in U is said to be **convergent** to $x \in U$ if given $t > 0, r \in (0, 1)$ there exist an integer $n_0 \in \mathbf{N}$ such that

$$N^*(x_n - x, t) < r \quad \forall n \geq n_0.$$

Definition 2.7 [1]. Let (U, N^*) be a fuzzy antinormed linear space. A sequence $\{x_n\}_n$ in U is said to be a **cauchy sequence** if given $t > 0$, $r \in (0, 1)$ there exist an integer $n_0 \in N$ such that

$$N^*(x_{n+p} - x_n, t) < r \quad \forall n \geq n_0, p = 1, 2, 3, \dots$$

Definition 2.8 [1]. A subset A of a fuzzy antinormed linear space (U, N^*) is said to be **bounded** if and only if there exist $t > 0$, $r \in (0, 1)$ such that

$$N^*(x, t) < r \quad \forall x \in A$$

Definition 2.9 [1]. A subset A of a fuzzy antinormed linear space (U, N^*) is said to be **compact** if for any sequence $\{x_n\}_n$ in A has a subsequence converging to an element of A .

Definition 2.10 [1]. Let (U, N^*) be a fuzzy antinormed linear space. A subset B of U is said to be **closed** if any sequence $\{x_n\}_n$ in B converges to $x \in B$, that is,

$$\lim_{n \rightarrow \infty} N^*(x_n - x, t) = 0, \quad \forall t > 0 \Rightarrow x \in B.$$

3 Fuzzy Anti-normed Linear Space

In this section we modify the definition of fuzzy antinorm and deduce some important theorems.

Definition 3.1 Let V be linear space over the field $F (= R \text{ or } C)$. A fuzzy subset ν of $V \times R$ is called a **fuzzy antinorm** on V if and only if for all $x, y \in V$ and $c \in F$

- (i) For all $t \in R$ with $t \leq 0$, $\nu(x, t) = 1$;
- (ii) For all $t \in R$ with $t > 0$, $\nu(x, t) = 0$ if and only if $x = \theta$;
- (iii) For all $t \in R$ with $t > 0$, $\nu(cx, t) = \nu(x, \frac{t}{|c|})$ if $c \neq 0, c \in F$;
- (iv) For all $s, t \in R$ with $\nu(x + y, s + t) \leq \nu(x, s) \diamond \nu(y, t)$;
- (v) $\lim_{t \rightarrow \infty} \nu(x, t) = 0$.

Definition 3.2 If $A^* = \{((x, t), \nu(x, t)) : (x, t) \in V \times R\}$ is a fuzzy antinorm on a linear space V over a field F . Then (V, A^*) is called a fuzzy antinormed linear space.

We further assume that for a fuzzy anti-normed linear space (V, A^*) ,

(vi) $\nu(x, t) < 1, \forall t > 0 \Rightarrow x = \theta$.

(vii) $\nu(x, \cdot)$ is a continuous function of $\mathbf{R}$ and strictly decreasing on the subset $\{t : 0 < \nu(x, t) < 1\}$ of $\mathbf{R}$.

Remark 3.3 Let ν be a fuzzy anti-norm on V then $\nu(x, t)$ is non-increasing with respect to t for each $x \in V$.

Proof. Let $t < s$. Then $k = s - t > 0$ we have

$$\nu(x, t) = \nu(x, t) \diamond 0 = \nu(x, t) \diamond \nu(0, k) \geq \nu(x, s).$$

Hence the proof.

Example 3.4 Let $(V, \|\cdot\|)$ be a normed linear space and define $a \diamond b = \max\{a, b\}$. For all $t(> 0) \in \mathbf{R}$, define

$$\nu(x, t) = \frac{k\|x\|}{t + k\|x\|}, \quad k > 0.$$

It is easy to verify that $A^* = \{((x, t), \nu(x, t)) : (x, t) \in V \times \mathbf{R}\}$ is a fuzzy anti-norm on V .

Example 3.5 Let $(V, \|\cdot\|)$ be a normed linear space, define $\nu : V \times \mathbf{R} \rightarrow [0, 1]$ by

$$\begin{aligned} \nu(x, t) &= 0, \text{ if } t > \|x\| \\ &= 1, \text{ if } t \leq \|x\| \end{aligned}$$

Then (V, ν) is a fuzzy anti-normed linear space.

Theorem 3.6 Let (V, A^*) be a fuzzy antinormed linear space. Then

$$\|x\|_\alpha^* = \wedge\{t : \nu(x, t) \leq 1 - \alpha\}, \quad \alpha \in (0, 1)$$

is an ascending family of α -norms on V .

Proof. Obvious.

Theorem 3.7 Let (V, A^*) be a fuzzy antinormed linear space satisfying (vi) and (vii). Let $\|x\|_\alpha^* = \wedge\{t : \nu(x, t) \leq 1 - \alpha\}, \alpha \in (0, 1)$. Also, let $\nu' : V \times \mathbf{R} \rightarrow [0, 1]$ be defined by

$$\begin{aligned} \nu'(x, t) &= \wedge\{1 - \alpha : \|x\|_\alpha^* \leq t\}, \text{ if } (x, t) \neq (\theta, 0) \\ &= 1, \text{ if } (x, t) = (\theta, 0) \end{aligned}$$

Then $\nu' = \nu$.

To prove this theorem we use the following lemmas.

Lemma 3.8 *Let (V, A^*) be a fuzzy antinormed linear space satisfying (vi) and (vii). Also, if $\{\|\cdot\|_\alpha^* : \alpha \in (0, 1)\}$ be ascending family of norms of V , defined by*

$$\|x\|_\alpha^* = \wedge \{t : \nu(x, t) \leq 1 - \alpha\}, \alpha \in (0, 1).$$

Then for $x_0 (\neq \theta) \in V$, $\nu(x_0, \|x_0\|_\alpha^) \leq 1 - \alpha, \forall \alpha \in (0, 1)$.*

Proof. Let $\|x_0\|_\alpha^* = s$, then $s > 0$. Then there exist a sequence $\{s_n\}_n$, $s_n > 0$ such that $\nu(x_0, s_n) \leq 1 - \alpha$, for all $n \in N$ and $s_n \rightarrow s$ as $n \rightarrow \infty$. Therefore

$$\begin{aligned} \lim_{n \rightarrow \infty} \nu(x_0, s_n) \leq 1 - \alpha &\Rightarrow \nu(x_0, \lim_{n \rightarrow \infty} s_n) \leq 1 - \alpha \quad \text{by (vii)} \\ &\Rightarrow \nu(x_0, \|x_0\|_\alpha^*) \leq 1 - \alpha, \forall \alpha \in (0, 1) \end{aligned}$$

Lemma 3.9 *Let (V, A^*) be a fuzzy antinormed linear space satisfying (vi) and (vii) and $\{\|\cdot\|_\alpha^* : \alpha \in (0, 1)\}$ be ascending family of norms of V , defined by*

$$\|x\|_\alpha^* = \wedge \{t : \nu(x, t) \leq 1 - \alpha\}, \alpha \in (0, 1).$$

Then for $x_0 (\neq \theta) \in V$, $\alpha \in (0, 1)$ and $s (> 0) \in R$,

$$\|x_0\|_\alpha^* = s \Leftrightarrow \nu(x_0, s) = 1 - \alpha$$

Proof. Let $\alpha \in (0, 1)$, $x (\neq \theta) \in V$ and $s = \|x_0\|_\alpha^* = \wedge \{t : \nu(x_0, t) \leq 1 - \alpha\}$. Now by lemma 3.9, we have

$$\nu(x_0, s) \leq 1 - \alpha \tag{1}$$

If possible let $\nu(x_0, s) < 1 - \alpha$ then by continuity of $\nu(x, \cdot)$ at s , there exist $s' < s$ such that $\nu(x_0, s') < 1 - \alpha$, which is impossible since $s = \wedge \{t : \nu(x_0, t) \leq 1 - \alpha\}$. Thus

$$\nu(x_0, s) \geq 1 - \alpha \tag{2}$$

From (1) and (2) it follows that $\nu(x_0, s) = 1 - \alpha$. Thus

$$\|x_0\|_\alpha^* = s \Rightarrow \nu(x_0, s) = 1 - \alpha \tag{3}$$

Next if $\nu(x_0, s) = 1 - \alpha$, $\alpha \in (0, 1)$ then from remark 3.3

$$\|x_0\|_\alpha^* = \wedge \{t : \nu(x_0, t) \leq 1 - \alpha\} = s. \tag{4}$$

Hence from (3) and (4) we have for $\alpha \in (0, 1)$, $x(\neq \theta) \in V$ and for $s > 0$, $\|x_0\|_\alpha^* = s \Leftrightarrow \nu(x_0, s) = 1 - \alpha$.

Proof of the main theorem:

Let $(x_0, t_0) \in V \times R$ and $\alpha \in (0, 1)$. To prove this theorem we consider the following cases:

case-1: For any $x_0 \in V$ and $t \leq 0$, $\nu(x_0, t_0) = \nu'(x_0, t_0) = 1$.

case-2: Let $x_0 = \theta$, $t_0 > 0$. Then $\nu(x_0, t_0) = \nu'(x_0, t_0) = 0$.

case-3: Let $x_0 \neq \theta$, $t_0 (> 0) \in R$ such that $\nu(x_0, t_0) = 1$. By lemma 3.1 we have, $\nu(x_0, \|x\|_\alpha^*) \leq 1 - \alpha$. Since $\nu(x_0, t_0) = 1 > 1 - \alpha$ it follows that $\nu(x_0, \|x\|_\alpha^*) \leq 1 - \alpha < \nu(x_0, t_0)$ and since $\nu(x_0, \cdot)$ is strictly non increasing $t_0 < \|x\|_\alpha^*$, $\forall \alpha(0, 1)$. So $\nu'(x_0, t_0) = \wedge \{1 - \alpha : \|x_0\|_\alpha^* \leq t_0\} = 1$. Thus $\nu(x_0, t_0) = \nu'(x_0, t_0) = 1$.

case-4: Let $x_0 \neq \theta$, $t_0 (> 0) \in R$ such that $\nu(x_0, t_0) = 0$. Note that,

$$\nu'(x, t) = \wedge \{1 - \alpha : \|x\|_\alpha^* \leq t\}, \text{ if } (x, t) \neq (\theta, 0) \quad (5)$$

$$\|x\|_\alpha^* = \wedge \{t : \nu(x, t) \leq 1 - \alpha\}, \alpha \in (0, 1], x \in V. \quad (6)$$

It follows that, $\|x_0\|_\alpha^* < t_0$, $\forall \alpha \in (0, 1)$, by (6). Therefore, $\|x_0\|_\alpha^* < t_0 \Rightarrow \nu'(x_0, t_0) = 0$, by (5). Thus, $\nu(x_0, t_0) = \nu'(x_0, t_0) = 0$.

case-5: Let $x_0 \neq \theta$, $t_0 (> 0) \in R$ such that $0 < \nu(x_0, t_0) < 1$. Note that $\forall \alpha \in (0, 1)$

$$\nu'(x, t) = \wedge \{1 - \alpha : \|x\|_\alpha^* \leq t\}, \text{ if } (x, t) \neq (\theta, 0) \quad (7)$$

$$\|x\|_\alpha^* = \wedge \{t : \nu(x, t) \leq 1 - \alpha\}, \alpha \in (0, 1], x \in V. \quad (8)$$

Let, $\nu(x_0, t_0) = 1 - \beta$, then from (8) we have

$$\|x\|_\beta^* \leq t_0. \quad (9)$$

Using (9) from (7) we get, $\nu'(x_0, t_0) \leq 1 - \beta$. Therefore,

$$\nu(x_0, t_0) \geq \nu'(x_0, t_0) \quad (10)$$

Now from lemma 3.2 we have $\nu(x_0, t_0) = 1 - \beta \Leftrightarrow \|x\|_\beta^* = t_0$.

Now, for $\beta < \alpha < 1$, let $\|x\|_\alpha^* = t'$. Then by lemma 3.2 we have $\nu(x_0, t') = 1 - \alpha$. So, $\nu(x_0, t') = 1 - \alpha < 1 - \beta = \nu(x_0, t_0)$. since, $\nu(x_0, \cdot)$ is strictly monotonically decreasing and $\nu(x_0, t') < \nu(x_0, t_0)$. Therefore $t' > t_0$. Then for $\beta < \alpha < 1$, we have $\|x\|_\alpha^* = t' > t_0$. So,

$$\nu'(x_0, t_0) \geq 1 - \beta = \nu(x_0, t_0) \quad (11)$$

Thus from (10) and (11) we have $\nu(x_0, t_0) = \nu'(x_0, t_0)$.

Since, $(x_0, t_0) \in V \times R$ is arbitrary $\nu'(x, t) = \nu(x, t_0)$, for all $(x, t) \in V \times R$.

Lemma 3.10 (Riesz): Let W be a closed and proper subspace of a fuzzy antinormed linear space (V, ν) satisfying (vi) and (vii). Then for each $\epsilon > 0$ there exist $y \in V - W$ such that $\nu(y, 1) \leq 1 - \alpha$ and $\nu(y - w, 1 - \epsilon) \leq 1 - \alpha$ for all $\alpha \in (0, 1)$ and $w \in W$.

Proof. Recall that, $\|x\|_\alpha^* = \bigwedge \{t : \nu(x, t) \leq 1 - \alpha\}$, $\alpha \in (0, 1)$ and $\{\|\cdot\|_\alpha^* : \alpha \in (0, 1)\}$ is an ascending family of α -norms on a linear space V . Now by applying Riesz lemma for normed linear space, it follows that for any $\epsilon > 0$ there exist $y \in V - W$ such that

$$\|y\|_\alpha^* = 1 \quad (12)$$

$$\|y - w\|_\alpha^* > 1 - \epsilon \quad \forall w \in W \quad (13)$$

Now, from theorem 3.7, for all $\alpha \in (0, 1)$ we have

$$\begin{aligned} \nu(y, t) &= \bigwedge \{1 - \alpha : \|y\|_\alpha^* \leq t\} \\ &\Rightarrow \nu(y, 1) = \bigwedge \{1 - \alpha : \|y\|_\alpha^* \leq 1\} \\ &\Rightarrow \nu(y, 1) \leq 1 - \alpha \end{aligned}$$

Again,

$$\begin{aligned} \nu(y - w, t) &= \bigwedge \{1 - \alpha : \|y - w\|_\alpha^* \leq t\} \\ &\Rightarrow \nu(y - w, \epsilon) = \bigwedge \{1 - \alpha : \|y - w\|_\alpha^* \leq \epsilon\} \\ &\Rightarrow \nu(y - w, \epsilon) \leq 1 - \alpha \end{aligned}$$

Hence the proof.

Theorem 3.11 Let (V, A^*) be a fuzzy antinormed linear space satisfying (vi) and (vii). If the set $\{x : \nu(x, 1) \leq 1 - \alpha\}$, $\alpha \in (0, 1)$ is compact then V is a space of finite dimension.

To prove this theorem we use the following lemma

Lemma 3.12 *In a fuzzy antinormed linear space (V, A^*) satisfying (vi) and (vii), every sequence is convergent if and only if it is convergent with respect its corresponding α -norms, $\alpha \in (0, 1)$*

Proof. $\Rightarrow$ *part* : Let (V, A^*) be a fuzzy antinormed linear space satisfying (vi) and (vii) and $\{x_n\}_n$ be a sequence in V such that $x_n \rightarrow x$.

$$\lim_{n \rightarrow \infty} \nu(x_n - x, t) = 0, \forall t > 0$$

Choose $0 < \alpha < 1$. So, $\lim_{n \rightarrow \infty} \nu(x_n - x, t) = 0 < 1 - \alpha \Rightarrow$ there exist $n_0(t)$ such that

$$\nu(x_n - x, t) < 1 - \alpha \quad \forall n \geq n_0(t, \alpha) \quad (14)$$

Now,

$$\|x_n - x\|_\alpha^* = \wedge \{t : \nu(x_n - x, t) \leq 1 - \alpha\} \Rightarrow \|x_n - x\|_\alpha^* \leq t, \forall n \geq n_0(t, \alpha)$$

by (3.3). Since, $t > 0$ is arbitrary,

$$\|x_n - x\|_\alpha^* \rightarrow 0 \text{ as } n \rightarrow \infty, \forall \alpha \in (0, 1)$$

$\Leftarrow$ *part* : Next, we suppose that, $\|x_n - x\|_\alpha^* \rightarrow 0$ as $n \rightarrow \infty, \forall \alpha \in (0, 1)$.

Then for $\alpha \in (0, 1), \epsilon > 0$ there exist $n_0(\alpha, \epsilon)$ such that

$$\|x_n - x\|_\alpha^* < \epsilon, \forall n \geq n_0(\alpha, \epsilon), \alpha \in (0, 1) \quad (15)$$

Now,

$$\nu(x_n - x, \epsilon) = \wedge \{1 - \alpha : \|x_n - x\|_\alpha^* \leq \epsilon\}$$

$$\Rightarrow \nu(x_n - x, \epsilon) \leq 1 - \alpha, \forall n \geq n_0(\alpha, \epsilon), \alpha \in (0, 1)$$

$$\Rightarrow \lim_{n \rightarrow \infty} \nu(x_n - x, \epsilon) = 0$$

Thus x_n converges to x .

Proof of the main theorem: It can be easily verified that $\{x : \nu(x, 1) \leq 1 - \alpha\} = \{x : \|x\|_\alpha^* \leq 1\}, \alpha \in (0, 1)$. By applying Riesz lemma, it can be proved that if for some $\alpha \in (0, 1)$ the set $\{x : \|x\|_\alpha^* \leq 1\}$ is compact then V is of finite dimensional. Using lemma 3.4, it follows that, for some $\alpha \in (0, 1), \{x : \nu(x, 1) \leq 1 - \alpha\}$ is compact then V of finite dimensional.

4 Fuzzy α -anti-convergence

In this section, the relations of fuzzy α -anti-convergence, fuzzy α -anti-cauchyness, fuzzy α -anti-compactness with their corresponding α -norms are studied and a few properties are derived.

Theorem 4.1 *Let (V, A^*) be a fuzzy antinormed linear space satisfying (vi) and (vii) and $\{\|\cdot\|_\alpha^* : \alpha \in (0, 1)\}$ be ascending family of norms of V , defined by*

$$\|x\|_\alpha^* = \wedge \{t : \nu(x, t) \leq 1 - \alpha\}, \alpha \in (0, 1).$$

Then for any increasing (or, decreasing) sequence $\{\alpha_n\}_n$ in $(0, 1)$, $\alpha_n \rightarrow \alpha$ in $(0, 1)$ implies $\|x\|_{\alpha_n}^ \rightarrow \|x\|_\alpha^*, \forall x \in V$*

Proof. For $x = \theta$, it is clear that α_n converges to $\alpha \Rightarrow \|x\|_{\alpha_n}^* \rightarrow \|x\|_\alpha^*$ suppose $x \neq \theta$. Then from Lemma 3.9, for $x \neq \theta, \alpha \in (0, 1), t' > 0$ we have

$$\|x\|_\alpha^* = t' \Leftrightarrow \nu(x, t') = 1 - \alpha$$

Let $\{\alpha_n\}_n$ be an increasing sequence in $(0, 1)$ such that α_n converges to α in $(0, 1)$.

Let $\|x\|_{\alpha_n}^* = s_n$ and $\|x\|_\alpha^* = s$

Then,

$$\nu(x, s_n) = 1 - \alpha_n \text{ and } \nu(x, s) = 1 - \alpha. \quad (16)$$

Since $\{\|\cdot\|_\alpha^* : \alpha \in (0, 1)\}$ is increasing family of norms and $\{s_n\}_n$ is increasing sequence of real numbers and bounded above by s . Hence $\{s_n\}_n$ is convergent. Thus,

$$\lim_{n \rightarrow \infty} \nu(x, s_n) = 1 - \lim_{n \rightarrow \infty} \alpha_n \Rightarrow \nu(x, \lim_{n \rightarrow \infty} s_n) = 1 - \alpha \quad (17)$$

From (5.1) and (5.2) we have, $\nu(x, \lim_{n \rightarrow \infty} s_n) = \nu(x, s)$ this implies $\lim_{n \rightarrow \infty} s_n = s$, by (vii). Therefore,

$$\lim_{n \rightarrow \infty} \|x\|_{\alpha_n}^* = \|x\|_\alpha^*$$

Similarly, if $\{\alpha_n\}_n$ is a decreasing sequence in $(0, 1)$ and α_n converges to α in $(0, 1)$ then it can be easily shown that $\|x\|_{\alpha_n}^* \rightarrow \|x\|_\alpha^*, \forall x \in V$.

Let V be a linear space and $\{\|\cdot\|_\alpha^* : \alpha \in (0, 1)\}$ be ascending family of fuzzy α -anti-norms on V . Assume that

(viii) for any increasing (or, decreasing) sequence $\{\alpha_n\}_n$ in $(0, 1)$, if α_n converges to α in $(0, 1)$, then $\|x\|_{\alpha_n}^*$ converges to $\|x\|_\alpha^*$.

Theorem 4.2 Let (V, A^*) be a fuzzy antinormed linear space and $\{\|\cdot\|_\alpha^* : \alpha \in (0, 1)\}$ be ascending family of norms of V satisfying (viii). Let, ν be fuzzy antinorm on V defined by

$$\begin{aligned} \nu(x, t) &= \wedge \{1 - \alpha : \|x\|_\alpha^* \leq t\}, \text{ if } (x, t) \neq (\theta, 0) \\ &= 1, \text{ if } (x, t) = (\theta, 0) \end{aligned} \quad (18)$$

Let $\|x\|'_\alpha : V \longrightarrow R^+$ be a function defined by

$$\|x\|'_\alpha = \wedge \{t : \nu(x, t) \leq 1 - \alpha\}. \quad (19)$$

Then $\|x\|'_\alpha = \|x\|_\alpha^*$, $\forall \alpha \in (0, 1)$.

Proof. For $x = \theta$, $\|x\|'_\alpha = \|x\|_\alpha^*$, $\forall \alpha \in (0, 1)$.
Consider when $x \neq \theta$. Let, $\alpha_0 \in (0, 1)$. Putting $\|x\|_\alpha^* = s_0$. Then $s_0 > 0$.
From (18) we get, $\nu(x, s_0) \leq \alpha_0$.
From (19) we get, $\|x\|'_{\alpha_0} \geq s_0 = \|x\|_{\alpha_0}^*$. Thus

$$\|x\|'_{\alpha_0} \geq \|x\|_{\alpha_0}^* \quad (20)$$

Next let $s > \|x\|'_{\alpha_0}$
 $\Rightarrow \exists s_1 < s$ such that $\nu(x, s_1) \leq \alpha_0$
 $\Rightarrow \wedge \{1 - \alpha : \|x\|_\alpha^* \leq \alpha_0\}$
If $\wedge \{1 - \alpha : \|x\|_\alpha^* = \alpha_0\}$, then there exist a decreasing sequence $\{\alpha_n\}_n$ in $(0, 1)$ such that $\alpha_n \longrightarrow \alpha_0$ and $\|x\|_{\alpha_n}^* \leq s_1$
Thus by (viii) we have $\|x\|_{\alpha_0}^* \leq s_1 < s$.
If $\wedge \{1 - \alpha : \|x\|_\alpha^* < \alpha_0\}$, then it follows that $\|x\|_{\alpha_n}^* \leq s_1 < s$.
Hence in any case $s > \|x\|_{\alpha_0}^*$. Thus

$$\|x\|'_{\alpha_0} \leq \|x\|_{\alpha_0}^* \quad (21)$$

Form (20) and (21) we get, $\|x\|'_{\alpha_0} = \|x\|_{\alpha_0}^*$
Since $\alpha_0 > 0$ is arbitrary, $\|x\|'_\alpha = \|x\|_\alpha^*$, $\forall x \in V$ and $\forall \alpha \in (0, 1)$,

Definition 4.3 Let (V, A^*) be a fuzzy antinormed linear space and $\alpha \in (0, 1)$. A sequence $\{x_n\}_n$ in V is said to be fuzzy α -anti-convergent in (V, A^*) if there exist $x \in V$ such that for all $t > 0$,

$$\lim_{n \rightarrow \infty} \nu(x_n - x, t) < 1 - \alpha$$

x is called fuzzy α -antilimit of x_n .

Theorem 4.4 *Let (V, A^*) be a fuzzy antinormed linear space satisfying (vi). Then fuzzy α -antilimit of a fuzzy α -anti-convergent sequence is unique.*

Proof. Let $\{x_n\}_n$ be a fuzzy α -anti-convergent sequence and suppose it converges to x and y in V . Then for all $t > 0$,

$$\lim_{n \rightarrow \infty} \nu(x_n - x, t) < 1 - \alpha \quad \text{and} \quad \lim_{n \rightarrow \infty} \nu(x_n - y, t) < 1 - \alpha.$$

Now,

$$\begin{aligned} \nu(x - y, t) &= \nu(x - x_n + x_n - y, t), \forall n. \\ &= \nu(x_n - x, t) \diamond \nu(x_n - y, t), \forall n. \end{aligned}$$

Taking limit we have

$$\begin{aligned} \nu(x - y, t) &= \lim_{n \rightarrow \infty} \nu(x_n - x, t) \diamond \lim_{n \rightarrow \infty} \nu(x_n - y, t) \\ &< (1 - \alpha) \diamond (1 - \alpha) = (1 - \alpha) \end{aligned}$$

That is, $\nu(x - y, t) < 1$ for all $t > 0$.

Therefore, $x - y = \theta$ by (vi)

$\Rightarrow x = y$.

Theorem 4.5 *Let (V, A^*) be a fuzzy antinormed linear space satisfying (vi). If $\{x_n\}_n$ be a fuzzy α -anti-convergent sequence in (V, A^*) such that x_n converges to x . Then $\|x_n - x\|_\alpha^* \rightarrow 0$ as $n \rightarrow \infty$.*

Proof. Since $\{x_n\}_n$ be a fuzzy α -anti-convergent sequence, suppose it converges to x , then for all $t > 0$,

$$\lim_{n \rightarrow \infty} \nu(x_n - x, t) < 1 - \alpha.$$

$$\Rightarrow \exists n_0(t) > 0 \text{ such that } \nu(x_n - x, t) < 1 - \alpha, \forall n \geq n_0(t).$$

$$\Rightarrow \exists n_0(t) > 0 \text{ such that } \|x_n - x\|_\alpha^* \leq t \quad \forall n \geq n_0(t).$$

Since $t > 0$ is arbitrary, $\|x_n - x\|_\alpha^* \rightarrow 0$ as $n \rightarrow \infty$.

Definition 4.6 *Let (V, A^*) be a fuzzy anti-normed linear space and $\alpha \in (0, 1)$. A sequence $\{x_n\}_n$ in V is said to be fuzzy α -anti-cauchy sequence if*

$$\lim_{n \rightarrow \infty} \nu(x_n - x_{n+p}, t) \leq 1 - \alpha \quad \forall t > 0, p = 1, 2, 3, \dots$$

Theorem 4.7 *Let (V, A^*) be a fuzzy antinormed linear space and $\alpha \in (0, 1)$. Then every fuzzy α -anti-convergent sequence in (V, A^*) is a fuzzy α -anti-cauchy sequence in (V, A^*) .*

Proof. Let $\{x_n\}_n$ be a fuzzy α -anti-convergent sequence and it converging to x . Then

$$\lim_{n \rightarrow \infty} \nu(x_n - x, t) < 1 - \alpha$$

Now,

$$\begin{aligned} \nu(x_n - x_{n+p}, t) &= \nu(x_n - x + x - x_{n+p}, t), \text{ for } p = 1, 2, 3, \dots \\ &= \nu(x_n - x, \frac{t}{2}) \diamond \nu(x_{n+p} - x, \frac{t}{2}), \text{ for } p = 1, 2, 3, \dots \end{aligned}$$

Therefore,

$$\begin{aligned} \lim_{n \rightarrow \infty} \nu(x_n - x_{n+p}, t) &\leq \lim_{n \rightarrow \infty} \nu(x_n - x, \frac{t}{2}) \diamond \lim_{n \rightarrow \infty} \nu(x_{n+p} - x, \frac{t}{2}) \\ &< (1 - \alpha) \diamond (1 - \alpha) = (1 - \alpha) \end{aligned}$$

Hence $\{x_n\}_n$ is a fuzzy α -anti-cauchy sequence in (V, A^*) .

Note 4.8 Every constant sequence in a fuzzy antinormed linear space (V, A^*) is a fuzzy α -anti-cauchy sequence in (V, A^*) , $\alpha \in (0, 1)$.

Proof. Obvious.

Theorem 4.9 Let (V, A^*) be a fuzzy antinormed linear space satisfying (vi). Then every Cauchy sequence in $(V, \|\cdot\|_\alpha^*)$ is a fuzzy α -anti-cauchy sequence in (V, A^*) , where $\|\cdot\|_\alpha^*$ denotes the α -norms on V , $\alpha \in (0, 1)$

Proof. Choose $\alpha_0 \in (0, 1)$ arbitrary but fixed. Let, $\{y_n\}_n$ be a Cauchy sequence in V with respect to $\|\cdot\|_{\alpha_0}^*$. Then

$$\lim_{n \rightarrow \infty} \|y_n - y_{n+p}\|_{\alpha_0}^* = 0.$$

Then for any given $\epsilon (> 0)$ there exist a positive integer $n_0(\epsilon)$ such that

$$\|y_n - y_{n+p}\|_{\alpha_0}^* < \epsilon, \forall n \geq n_0(\epsilon) \text{ and } p = 1, 2, 3, \dots$$

$$\Rightarrow \wedge \{t > 0 : \nu(y_n - y_{n+p}, t) \leq 1 - \alpha_0\} < \epsilon.$$

$$\Rightarrow \text{there exist } t(n, p, \epsilon) < \epsilon \text{ such that}$$

$$\nu(y_n - y_{n+p}, t(n, p, \epsilon)) \leq 1 - \alpha_0, \forall n \geq n_0(\epsilon) \text{ and } p = 1, 2, 3, \dots$$

$$\Rightarrow \nu(y_n - y_{n+p}, \epsilon) \leq 1 - \alpha_0$$

Since $\epsilon (> 0)$ is arbitrary,

$$\lim_{n \rightarrow \infty} \nu(y_n - y_{n+p}, t) \leq 1 - \alpha_0, \forall t > 0$$

$\Rightarrow \{y_n\}_n$ is fuzzy α_0 -anti-cauchy sequence in (V, A^*) .

Since, $\alpha_0 \in (0, 1)$ is arbitrary, every Cauchy sequence in (V, A^*) is fuzzy α -anti-cauchy sequence in (V, A^*) for each $\alpha \in (0, 1)$.

Definition 4.10 Let (V, A^*) be a fuzzy antinormed linear space and $\alpha \in (0, 1)$. It is said to be fuzzy α -anti-complete if every fuzzy α -anti-cauchy sequence in V fuzzy α -anti-converges to a point of V .

Theorem 4.11 Let (V, A^*) be a fuzzy antinormed linear space satisfying (vi). If (V, A^*) be fuzzy α -anti-complete then V be complete with respect to $\|\cdot\|_\alpha^*$, $\alpha \in (0, 1)$

Proof. Choose $\alpha_0 \in (0, 1)$ arbitrary but fixed. Let, $\{y_n\}_n$ be a cauchy sequence in V with respect to $\|\cdot\|_{\alpha_0}^*$. then it follows that $\{y_n\}_n$ is fuzzy α_0 -anti-cauchy sequence in (V, A^*) .

Since, (V, A^*) is fuzzy α_0 -anti-complete, there exist $y \in V$ such that

$$\begin{aligned} \lim_{n \rightarrow \infty} \nu(y_n - y, t) &< 1 - \alpha_0, \forall t > 0 \\ \Rightarrow \lim_{n \rightarrow \infty} \|y_n - y\|_{\alpha_0}^* &= 0, \text{ by theorem 4.5.} \\ \Rightarrow y_n &\longrightarrow y \text{ with respect to } \|\cdot\|_\alpha^*. \\ \Rightarrow (V, \|\cdot\|_{\alpha_0}^*) &\text{ is fuzzy anti-complete.} \end{aligned}$$

Since, α_0 is arbitrary, $(V, \|\cdot\|_\alpha^*)$ are complete.

5 Future Work

In our future work, our goal is to develop the concepts of different types of fuzzy anti-continuities and fuzzy anti-boundedness of linear operator and the relations among them.

References

- [1] Iqbal H. Jebril and T.K. Samanta, *Fuzzy anti-normed linear space*, Journal of mathematics and Technology, February, 2010.
- [2] T. Bag and S.K. Samanta, *Finite Dimensional Fuzzy Normed Linear Spaces*, The Journal of Fuzzy Mathematics Vol. 11 (2003) 687 - 705.
- [3] T. Bag and S.K. Samanta, *Fuzzy bounded linear operators*, Fuzzy Sets and Systems 151 (2005) 513 - 547.
- [4] S.C. Cheng and J.N. Mordeson, *Fuzzy Linear Operators and Fuzzy Normed Linear Spaces*, Bull. Cal. Math. Soc.86 (1994) 429 - 436.
- [5] Bivas Dinda and T.K. Samanta, *Intuitionistic Fuzzy Continuity and Uniform Convergence*, Int. J. Open Problems Compt.Math., Vol 3, No. 1 (2010) 8 - 26.

- [6] C. Felbin , *The completion of fuzzy normed linear space*, Journal of mathematical analysis and application 174(2) (1993) 428-440.
- [7] T.K. Samanta and Iqbal H. Jebril , *Finite dimentional intuitionistic fuzzy normed linear space*, Int. J. Open Problems Compt. Math., Vol 2, No. 4 (2009) 574-591.
- [8] A.K. Katsaras , *Fuzzy topological vector space*, Fuzzy Sets and Systems 12 (1984) 143 - 154.
- [9] S. Vijayabalaji, N. Thillaigovindan, Y.B. Jun *Intuitionistic Fuzzy n-normed linear space* , Bull. Korean Math. Soc. 44 (2007) 291 - 308.
- [10] O. Kramosil, J. Michalek , *Fuzzy metric and statisticalmetric spaces*, Kybernetika 11 (1975) 326 - 334.
- [11] B. Schweizer , A. Sklar, *Statistical metric space*, Pacific journal of mathematics 10 (1960) 314-334.
- [12] L.A. Zadeh *Fuzzy sets*, Information and control 8 (1965) 338-353.